

Images of a Bose-Einstein condensate at finite temperature *

Jacek Dziarmaga

*Institute of Physics and Centre for Complex Systems, Jagiellonian University, Reymonta 4, 30-059 Kraków, Poland
(June 28, 2005)*

A condensate initially prepared at finite temperature evolves under external time-dependent perturbation into a time-dependent mixed state. In these notes I use number-conserving time-dependent Bogoliubov theory to derive probability distribution for different outcomes of density measurement on the time-dependent excited state.

I. INTRODUCTION

Quantum measurements on Bose-condensed systems can give quite unexpected results. For example, in the classic paper by Javanainen and Yoo [1] a density measurement on a Fock state $|N/2, N/2\rangle$ with N particles equally divided between two counter-propagating plane waves $e^{\pm ix}$ reveals an interference pattern $\rho(x|\varphi) \sim \cos^2(x - \varphi)$ with a phase φ chosen randomly in every realization of the experiment. The Fock state has a uniform single particle density distribution, but its measurement unexpectedly reveals interference between the two counter-propagating condensates. The Fock state is a quantum superposition over N -particle condensates with different relative phases φ in their wave functions [2], $|N/2, N/2\rangle \sim \int d\varphi |N : e^{+i(x-\varphi)} + e^{-i(x-\varphi)}\rangle$, but every single realization of the experiment reveals such a density distribution as if the state before the density measurement were one of the condensates $|N : e^{+i(x-\varphi)} + e^{-i(x-\varphi)}\rangle$ with a randomly chosen phase φ . This effect is best explained [2] when the density measurement, which is a destructive measurement of all particle positions at the same time, is replaced by an equivalent sequential measurement of one position after another. With an increasing number n of measured positions a quantum state of the remaining $N - n$ particles gradually “collapses” from the initial uniform superposition over all phases to a state with a more and more localized phase φ . For a large N a measurement of only a small fraction $\frac{n}{N} \ll 1$ of all particles practically collapses the state of remaining $N - n$ particles to a condensate with definite phase φ .

A lesson from this instructive example [1,2] is that quantum measurement on an N -particle state with highly occupied single particle modes “collapses” the state to a definite condensate with a definite condensate wave function $\phi(x)$. The question is: what is the probability distribution for different measurement outcomes $\phi(x)$? As the set of condensates is not an orthonormal basis this is not a trivial question.

In Ref. [6] we derived this probability distribution in the framework of the time-dependent Bogoliubov theory at zero temperature. At zero temperature a condensate initially prepared in its N -particle ground state evolves under external time-dependent perturbation into a time-dependent excited state. The excited state is a time-dependent Bogoliubov vacuum i.e. at any time t there exists a complete set of quasiparticle annihilation operators for which the excited state is a vacuum. In Ref. [6] it was shown that the time-dependent vacuum has a simple diagonal structure which directly leads to a compact gaussian probability distribution for different condensate wave functions $\phi(x)$. As the case of zero temperature is covered in Ref. [6], in these notes I describe the general case of finite temperature when the initial state is a condensate in equilibrium with a thermal cloud of atoms. I derive gaussian probability distribution for $\phi(x)$ at any time t when the external perturbation drives the initial thermal state into an excited mixed state.

II. N-CONSERVING BOGOLIUBOV THEORY

Number conserving Bogoliubov theory [7] is a quadratic approximation to the second quantized Hamiltonian (in trap units)

$$\hat{H} = \int dx \left[\frac{1}{2} \partial_x \hat{\Psi}^\dagger \partial_x \hat{\Psi} + \frac{1}{2} x^2 \hat{\Psi}^\dagger \hat{\Psi} + V(t, x) \hat{\Psi}^\dagger \hat{\Psi} + \frac{1}{2} g \hat{\Psi}^\dagger \hat{\Psi}^\dagger \hat{\Psi} \hat{\Psi} \right]. \quad (1)$$

*Invited talk given at the conference “Quantum Optics VI”, June 13-18 2005, Krynica, Poland

Here $\hat{\Psi}(x)$ is the bosonic annihilation operator, $V(t, x)$ is the external perturbation potential, and g is strength of contact interaction between atoms. Here and in the following I will use one-dimensional notation but all equations can be generalized by the simple replacement $x \rightarrow \vec{x}$. The annihilation operator is split into condensate and non-condensate part

$$\hat{\Psi}(x) = \hat{a}_0 \phi_0(x) + \delta\hat{\psi}(x). \quad (2)$$

It is assumed that most atoms occupy the condensate mode $\phi_0(x)$. Equation (2) is substituted to the Hamiltonian (1) and then the Hamiltonian is expanded in powers of the fluctuation operator $\delta\hat{\psi}$, see Ref. [7].

Many experiments on dilute atomic condensates can be clearly divided in two steps: as a first step a condensate is prepared in its ground state and then in the second step an external potential $V(t, x)$ is applied to manipulate with the condensate wave function. Generic examples are phase imprinting of dark solitons [5], atomic interferometry [3], or generation of shock waves in Bose-Einstein condensates [4]. At finite temperature the initial state before the manipulation is a thermal state including thermal excitations above the N -particle ground state. The initial condensate wave function ϕ_0 solves the stationary Gross-Pitaevskii equation

$$\mu\phi_0 = -\frac{1}{2}\partial_x^2\phi_0 + \frac{1}{2}x^2\phi_0 + g|\phi_0|^2\phi_0. \quad (3)$$

In Bogoliubov approximation the ground state is Bogoliubov vacuum $|0_b\rangle$ which can be written as a gaussian superposition over condensates [8]

$$|0_b\rangle = \int d^2b e^{-\frac{1}{2}\sum_{m=1}^M b_m^* b_m} \left| N : \phi_0(x) + \frac{1}{\sqrt{N}} \sum_{m=1}^M b_m u_m(x) + b_m^* v_m^*(x) \right\rangle. \quad (4)$$

Here the state $|N : \phi\rangle$ is a condensate of N atoms in the normalized condensate wave function $\frac{\phi}{\sqrt{\langle\phi|\phi\rangle}}$. The Bogoliubov modes u_m and v_m are eigenmodes of the stationary Bogoliubov-de Gennes equations

$$\begin{aligned} \omega_m u_m &= -\frac{1}{2}\partial_x^2 u_m + \frac{1}{2}x^2 u_m + 2g|\phi_0|^2 u_m + g\phi_0^2 v_m, \\ -\omega_m v_m &= -\frac{1}{2}\partial_x^2 v_m + \frac{1}{2}x^2 v_m + 2g|\phi_0|^2 v_m + g(\phi_0^*)^2 u_m. \end{aligned} \quad (5)$$

Numerical solution of these equations gives a finite number of modes M . At finite temperature the initial state is a thermal state $\hat{\rho}(0)$ with thermal quasiparticle excitations. The thermal state is also a gaussian state [8]

$$\begin{aligned} \hat{\rho}(0) &= \int d^2b_L \int d^2b_R e^{-\frac{1}{2}b_L^* b_L - \frac{1}{2}b_R^* b_R + b_L^* e^{-\beta\omega} b_R} \\ &\quad \left| N : \phi_0(x) + \frac{1}{\sqrt{N}} \sum_{m=1}^M b_{L,m} u_m(x) + b_{L,m}^* v_m^*(x) \right\rangle \left\langle N : \phi_0(x) + \frac{1}{\sqrt{N}} \sum_{m=1}^M b_{R,m} u_m(x) + b_{R,m}^* v_m^*(x) \right|. \end{aligned} \quad (6)$$

Here $b_L^* b_L = \sum_{m=1}^M b_{L,m}^* b_{L,m}$ and $b_L^* e^{-\beta\omega} b_R = \sum_{m=1}^M b_{L,m}^* e^{-\beta\omega_m} b_{R,m}$.

In Bogoliubov theory the initial thermal state evolves under external perturbation $V(t, x)$ into an excited state $\hat{\rho}(t)$ which has the same form as the initial $\hat{\rho}(0)$ in Eq.(6) but with time-dependent Bogoliubov modes $u_m(t, x)$ and $v_m(t, x)$ which solve time-dependent Bogoliubov-de Gennes equations

$$\begin{aligned} i\partial_t u_m &= -\frac{1}{2}\partial_x^2 u_m + \frac{1}{2}x^2 u_m + 2g|\phi_0|^2 u_m + g\phi_0^2 v_m, \\ -i\partial_t v_m &= -\frac{1}{2}\partial_x^2 v_m + \frac{1}{2}x^2 v_m + 2g|\phi_0|^2 v_m + g(\phi_0^*)^2 u_m \end{aligned} \quad (7)$$

with initial conditions being the eigenmodes of the stationary BdG equations (5). The time-dependent condensate wave function $\phi_0(t, x)$ solves the time-dependent Gross-Pitaevskii equation

$$i\partial_t \phi_0 = -\frac{1}{2}\partial_x^2 \phi_0 + \frac{1}{2}x^2 \phi_0 + g|\phi_0|^2 \phi_0. \quad (8)$$

III. PROBABILITY DISTRIBUTION FOR OUTCOMES

As mentioned before, density measurement is “collapsing” N -particle state to a Bose-Einstein condensate. The aim of the measurement theory is to provide probability distribution for different condensate wave functions ϕ . In the present context of Bogoliubov theory it is convenient to split possible condensate wave functions into the condensate part and the non-condensate part: $\phi = \phi_0 + \frac{1}{\sqrt{N}}\delta\phi$. The aim is to find gaussian probability distribution for $\delta\phi$ in the gaussian state $\hat{\rho}(t)$. Ideally the gaussian distribution would be fully determined by the following equalities between second order correlators of the gaussian $\delta\phi(x)$ and second order correlators of the field operators:

$$\overline{\delta\phi^*(x)\delta\phi(y)} \stackrel{?}{=} \langle \delta\hat{\psi}^\dagger(x)\delta\hat{\psi}(y) \rangle = \sum_m n_m u_m^*(x)u_m(y) + (1+n_m)v_m(x)v_m^*(y) , \quad (9)$$

$$\overline{\delta\phi(x)\delta\phi^*(y)} \stackrel{?}{=} \langle \delta\hat{\psi}(x)\delta\hat{\psi}^\dagger(y) \rangle = \sum_m n_m v_m^*(x)v_m(y) + (1+n_m)u_m(x)u_m^*(y) , \quad (10)$$

$$\overline{\delta\phi(x)\delta\phi(y)} = \langle \delta\hat{\psi}(x)\delta\hat{\psi}(y) \rangle = \sum_m n_m v_m^*(x)u_m(y) + (1+n_m)u_m(x)v_m^*(y) , \quad (11)$$

$$\overline{\delta\phi^*(x)\delta\phi^*(y)} = \langle \delta\hat{\psi}^\dagger(x)\delta\hat{\psi}^\dagger(y) \rangle = \sum_m n_m u_m^*(x)v_m(y) + (1+n_m)v_m(x)u_m^*(y) . \quad (12)$$

Here the most right hand sides follow from the Bogoliubov theory [7]. $n_m = (e^{\beta\omega_m} - 1)^{-1}$ is average number of thermally excited Bogoliubov quasiparticles in the initial state. Unfortunately, because of the non-zero commutator $[\delta\hat{\psi}(x), \delta\hat{\psi}^\dagger(y)] = \delta(x-y) - \phi_0^*(x)\phi_0(y)$, the first two conditions cannot be satisfied simultaneously. I replace them with the condition

$$\overline{\delta\phi^*(x)\delta\phi(y)} = \left(\overline{\delta\phi(x)\delta\phi^*(y)} \right)^* = \frac{1}{2} \langle \delta\hat{\psi}^\dagger(x)\delta\hat{\psi}(y) + \delta\hat{\psi}(y)\delta\hat{\psi}^\dagger(x) \rangle . \quad (13)$$

It is convenient to expand the fluctuation as

$$\delta\phi(x|z) = \sum_{\alpha=1}^{\infty} z_{\alpha} \phi_{\alpha}(x) \quad (14)$$

in the ortonormal basis of the eigenmodes ϕ_{α} of the reduced single particle density matrix

$$\langle \delta\hat{\psi}^\dagger(x)\delta\hat{\psi}(y) \rangle = \sum_{\alpha=1}^M \delta N_{\alpha} \phi_{\alpha}^*(x)\phi_{\alpha}(y) . \quad (15)$$

Here the left hand side is given by Eq.(9). The right hand side is obtained after diagonalization of the hermitean operator on the left. The real eigenvalues δN_{α} are average occupation numbers of the corresponding non-condensate modes ϕ_{α} . The correlators (12) after the semiclassical approximation (13) determine the matrix of correlators of the complex gaussian random variables z_{α} :

$$\begin{pmatrix} \overline{z_{\alpha}^* z_{\beta}} & \overline{z_{\alpha}^* z_{\beta}^*} \\ \overline{z_{\alpha} z_{\beta}} & \overline{z_{\alpha} z_{\beta}^*} \end{pmatrix} = \begin{pmatrix} D_{\alpha\beta} & C_{\alpha\beta} \\ C_{\alpha\beta}^* & D_{\alpha\beta} \end{pmatrix} . \quad (16)$$

Here the $M \times M$ matrices on the right hand side are

$$D_{\alpha\beta} = \frac{1}{2} (U_{\alpha m}^* n_m U_{\beta m} + V_{\alpha m}^* (1+n_m) V_{\beta m} + U_{\alpha m} (1+n_m) U_{\beta m}^* + V_{\alpha m} n_m V_{\beta m}^*) , \quad (17)$$

$$C_{\alpha\beta} = U_{\alpha m}^* n_m V_{\beta m}^* + V_{\alpha m}^* (1+n_m) U_{\beta m}^* , \quad (18)$$

with the matrix elements $U_{\alpha m} = \langle \phi_{\alpha} | u_m \rangle$ and $V_{\alpha m} = \langle \phi_{\alpha} | v_m^* \rangle$. Replacing z_{α} 's with real coordinates, $z_{\alpha} = x_{\alpha} + iy_{\alpha}$, we get a real symmetric matrix of correlators

$$\begin{pmatrix} \overline{x_{\alpha} x_{\beta}} & \overline{x_{\alpha} y_{\beta}} \\ \overline{y_{\alpha} x_{\beta}} & \overline{y_{\alpha} y_{\beta}} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \text{Re } D_{\alpha\beta} + \text{Re } C_{\alpha\beta} & \text{Im } C_{\alpha\beta} \\ \text{Im } C_{\alpha\beta} & \text{Re } D_{\alpha\beta} - \text{Re } C_{\alpha\beta} \end{pmatrix} \quad (19)$$

and a condition that $\text{Im } D_{\alpha\beta} = 0$ - a good test of the correctness of the calculations. Diagonalization of the correlation matrix (19) gives eigenvalues $\lambda_s \geq 0$ with $s = 1, \dots, 2M$. Corresponding eigenvectors are columns of an orthogonal matrix O . The eigenvectors define convenient parametrization of the gaussian fluctuation as

$$\delta\phi(x) = \sum_{\alpha=1}^M z_{\alpha} \phi_{\alpha}(x) = \sum_{\alpha=1}^M \phi_{\alpha}(x) \sum_{s=1}^{2M} (O_{\alpha,s} + iO_{M+\alpha,s}) q_s \equiv \sum_{s=1}^{2M} \Phi_s(x) q_s \quad (20)$$

with independent real gaussian random variables q_s of zero mean and variances $\overline{q_s^2} = \lambda_s$. However, this is not the end of the story yet.

As a result of the semiclassical approximation in Eq.(13) averages like e.g. average density of depletion $\overline{\delta\phi^*(x)\delta\phi(x)}$ are divergent because there is infinite number of unoccupied modes $\phi_{\alpha}(x)$, every one of them contributing to this depletion density a term $\frac{1}{2}\phi_{\alpha}^*(x)\phi_{\alpha}(x)$. In stochastic averages like $\overline{\delta\phi^*(x)\delta\phi(x)}$ average occupation numbers of modes $\phi_{\alpha}(x)$ seem to be $\delta N_{\alpha} + \frac{1}{2}$ instead of the correct δN_{α} . This artifact of the semiclassical approximation can be corrected by introducing to Eq.(20) of regularization factors:

$$\delta\phi(x)|_{\text{reg}} = \sum_{\alpha=1}^M z_{\alpha} \left(\frac{\delta N_{\alpha}}{\delta N_{\alpha} + \frac{1}{2}} \right)^{1/2} \phi_{\alpha}(x) = \sum_{\alpha=1}^M \left(\frac{\delta N_{\alpha}}{\delta N_{\alpha} + \frac{1}{2}} \right)^{1/2} \phi_{\alpha}(x) \sum_{s=1}^{2M} (O_{\alpha,s} + iO_{M+\alpha,s}) q_s \equiv \sum_{s=1}^{2M} \Phi_s(x) q_s \quad (21)$$

As expected in semiclassical approximation, the regularizing factors $\left(\frac{\delta N_{\alpha}}{\delta N_{\alpha} + \frac{1}{2}} \right)^{1/2}$ are approximately 1 for the highly occupied modes with $\delta N_{\alpha} \gg 1$ which dominate in the density distribution, but at the same time they remove the divergence coming from the infinity of unoccupied modes. The wave functions $\Phi_s(x)$ are in general neither normalized nor orthogonal, except in the quantum limit of zero temperature, see the proof in Ref. [6].

IV. CONCLUSION

In conclusion, a recipe to simulate density measurement on the time-dependent excited thermal state has the following steps:

- Solve stationary Gross-Pitaevskii and Bogoliubov-de Gennes equations (3,5) to provide initial conditions for $\phi_0(t, x)$, $u_m(t, x)$ and $v_m(t, x)$, and the initial quasiparticle frequencies ω_m .
- Solve time-dependent Gross-Pitaevskii and Bogoliubov-de Gennes equations (8,7) with respect to $\phi_0(t, x)$, $u_m(t, x)$, and $v_m(t, x)$.
- Diagonalize the reduced single particle matrix (15) to get its non-condensate eigenmodes ϕ_{α} with their average occupation numbers δN_{α} .
- Build the matrices $D_{\alpha\beta}$ and $C_{\alpha\beta}$ in Eqs.(17,18), and then the real symmetric correlation matrix in Eq.(19).
- Diagonalize the correlation matrix in Eq.(19) to get its real eigenvalues λ_s and corresponding eigenvectors $O_{\alpha s}$.
- Build the regularized modes Φ_s according to their definition implicit in Eq.(21):

$$\Phi_s(x) = \sum_{\alpha=1}^M \left(\frac{\delta N_{\alpha}}{\delta N_{\alpha} + \frac{1}{2}} \right)^{1/2} \phi_{\alpha}(x) (O_{\alpha,s} + iO_{M+\alpha,s}) . \quad (22)$$

- Choose independent real random variables q_s 's from their gaussian distributions of zero mean and variance $\overline{q_s^2} = \lambda_s$, and then combine the chosen q 's into condensate density

$$\rho(x|q) = \left| \sqrt{N}\phi_0(x) + \sum_{s=1}^{2M} q_s \Phi_s(x) \right|^2 . \quad (23)$$

The $\rho(x|q)$ defines a family of all possible density measurement outcomes with a gaussian probability distribution for different q 's.

In the limit of zero temperature this general recipe coincides with the recipe derived by different methods in Ref. [6]. At zero temperature the wave functions become $\Phi_{\alpha}(x) \sim \phi_{\alpha}(x)$ for $\alpha = 1, \dots, M$ and zero otherwise (here the $\sim$ means equality up to a phase factor). Corresponding variances are $q_{\alpha}^2 = \delta N_{\alpha}$.

ACKNOWLEDGEMENTS

I would like to thank Zbyszek Karkuszewski and Krzysztof Sacha for stimulating discussions. This work was supported in part by Polish government scientific funds (2005-2008) as a research project.

- [1] J. Javanainen and S.Mi. Yoo, Phys. Rev. Lett. **76**, 161 (1996).
- [2] Y. Castin and J. Dalibard, Phys. Rev. A **55**, 4330 (1997); K. Mølmer, Phys. Rev. A **65**, 021607 (2002); S. Ashhab and A.J. Leggett, Phys. Rev. A **65**, 023604 (2002).
- [3] Y. Shin *et al.*, Phys. Rev. Lett. **92**, 050405 (2004).
- [4] B. Damski, Phys. Rev. A **69**, 043610 (2004).
- [5] S. Burger *et al.*, Phys. Rev. Lett. **83**, 5198 (1999).
- [6] J. Dziarmaga and K. Sacha, cond-mat/0503328.
- [7] M. D. Girardeau and R. Arnowitt, Phys. Rev. **113**, 755 (1959); C. W. Gardiner, Phys. Rev. A **56**, 1414 (1997); M. D. Girardeau Phys. Rev. A **58**, 775 (1998); Y. Castin and R. Dum, Phys. Rev. A **57**, 3008 (1998); J. Dziarmaga and K. Sacha, Phys. Rev. A **67** 033608 (2003).
- [8] J. Dziarmaga and J. Meisner, cond-mat/0410707.